Conducted by: Chris Nagle - Tract #23

George Irwin -Tract #22
David McAndrew - Tract #5

White-Tailed Deer Spotlight Survey - 2009

Count	Date	Acres Visibile	Buck	Doe	Fawn	Unidentified	Total Observed	Density - Acres/Deer
1	9/20/2009	75	2	35	8	8	53	1.42
2	9/27/2009	75	2	30	8	2	42	1.79
3	10/4/2009	75	2	23	10	9	44	1.70
4	10/10/2009	75	9	34	4	5	52	1.44
Total		300	15	122	30	24	191	1.59

Cumulative Census Statistics:

1.59 Acres	per Deer							
8.13								
0.25								
Percentage of Deer Population by Category								
7.85%								
63.87%								
15.71%								
nch	523							
41.08								
334.08								
82.15								
	75							
8								
67								
1.85 Acres	per Deer							
	0.25 Category 7.85% 63.87% 15.71% nch 41.08 334.08 82.15							

Harvest Recommendations:

- 1) Heavy emphasis must be placed on the harvesting of Doe. Over the last several years the reanch has harvested fewer doe then harvest estimates and as a result, the population is approaching overpopulation.
- 2) The harvest of spike bucks should be included in the buck harvest quota, not added to it. Harvesting spike buck will remove poor quality buck from the herd at an early age.
- 3) The harvest of young, immature buck should be restricted. Deer body characteristics, in addition to antler characteristics, should be used to determine the relative age of buck on-the-hoof.

Records Management:

As set forth by TPWD, the following information will be included:

- A) Sex
- B) Weight
- C) Antler measurements; Inside spread & number of points
- D) Age